
References

- 1 Shapiro LJ: Steroid sulfatase deficiency and the genetics of the short arm of the human X chromosome. *Adv Hum Genet* 1985;14:331-381.
- 2 David M, Israel N, Merksamer R, et al: Very low maternal serum unconjugated estriol and prenatal diagnosis of steroid sulfatase deficiency. *Fetal Diagn Ther* 1995;10:76-80.
- 3 Lebo RV, Lynch ED, Golbus MS, et al: Prenatal in situ hybridization test for deleted steroid sulfatase gene. *Am J Med Genet* 1993;46:652-658.
- 4 Santolaya-Forgas J, Burd LI, Burton BK: Clinical significance of low levels of second trimester maternal serum human chorionic gonadotropin. *Fetal Diagn Ther* 1994;9:362-366.
- 5 Santolaya-Forgas J, Jessup J, Burd LI, Prins GS, Burton BK: Pregnancy outcome in women with low midtrimester maternal serum unconjugated estriol. *J Reprod Med* 1996;41:87-90.
- 6 Novice FM, Collison DW, Burgdorf WHC, Esteryl NB: Disorders of cornification: in: *Manual Book of Genetic Skin Disorders*. Philadelphia, Saunders, 1994, pp 1-85.

Announcement

14th International Congress: The Fetus as a Patient

Amsterdam, May 12-16, 1998

Venue: RAI Congress Centre.

Organization: The International Society The Fetus as a Patient.

Chairman:

Prof. Dr. H.P. van Geijn, chairman of the local organizing committee
Department of Obstetrics and Gynecology
University Hospital, Vrije Universiteit
NL-Amsterdam (The Netherlands)

Tel. +31 20 444 22 27, Fax +31 20 444 48 11

Information:

Congress Office SCEM
Ms. M. de Haan
PO Box 21
NL-4196 ZG Tricht (The Netherlands)

Tel. +31 345 57 66 42, Fax +31 345 57 17 81, E-mail SCEM@SCEM.NL