

Nutrition as a Potential Factor of Primary Dysmenorrhea: A Systematic Review of Observational Studies

Zahra Bajalan^a Zainab Alimoradi^a Farnoosh Moafi^b

^aSocial Determinants of Health Research Center, Qazvin University of Medical Sciences, Qazvin, Iran;

^bDepartment of Midwifery, School of Nursing and Midwifery, Qazvin University of Medical Sciences, Qazvin, Iran

Keywords

Nutrition · Primary dysmenorrhea · Systematic review · Observational studies

Abstract

Background: The incidence and severity of primary dysmenorrhea are influenced by various factors. The aim of the present study was to review nutritional factors influencing primary dysmenorrhea. **Methods:** Academic databases including Web of Science, EMBASE, Scopus, and PubMed (including Medline) were searched using keywords of nutrition, diet, and primary dysmenorrhea. In this study, observational studies that were published in English from 1990 to April 2018, which focused on nutritional factors affecting primary dysmenorrhea, were selected. The evaluation of studies was performed using a modified STROBE checklist with 10 items. **Results:** Out of 5,814 retrieved studies, 38 articles met inclusion criteria and were included for final data synthesis. The increased consumption of fruits and vegetables as the sources of vitamins and minerals, as well as fish and milk and dairy products have positive associations with less menstrual pain. Inconsistent results were reported on the consumption

of other nutritional groups. Studies showed negative associations of meal skipping and following diet to lose weight with severity of dysmenorrhea. **Conclusion:** A few studies showed inconclusive findings due to methodological heterogeneities for assessing nutritional habits and different methods of measuring dysmenorrhea pain. Therefore, further analysis and future interventional studies with stronger methodologies are required.

© 2019 S. Karger AG, Basel

Introduction

Dysmenorrhea is the most common health-related problem (60%) in women in the reproductive age [1, 2]. It can reduce the quality of life and hinders social activities in young women, especially when it is accompanied by symptoms such as headache, fatigue, nausea and vomiting, diarrhea, boredom, chills and muscle cramps [3–5]. Primary dysmenorrhea is responsible for disability and inefficiency in terms of absence from the school or workplace. It has been reported that about 1% of women in the reproductive age lose their active work hours for 1–3 days

per month. Also, about 14% of girls are absent from the school due to painful uterus contractions [6, 7]. The feelings of shame and mood changes due to dysmenorrhea influence women's social relationships [8]. Also, the consequences and economic losses of medications may happen [8]. Therefore, dysmenorrhea is considered important in terms of its economic and social effect on women.

The increased production or unbalanced levels of prostaglandins creates pain due to increased uterine contractility, decreased uterine blood flow, and increased sensitivity of peripheral nerves [2, 7]. Age, family history, body mass index, socioeconomic status, education, smoking, and alcohol use can influence the severity of primary dysmenorrhea [1, 7–17]. Non-steroidal anti-inflammatory drugs and oral contraceptive pills have been listed as common treatments for women with dysmenorrhea [18]. The goal of treatment is to provide adequate pain relief and reduction of symptoms with least adverse effects. Marjoribanks et al. [19] in a systematic review reported that non-steroidal anti-inflammatory drugs were more effective than placebos, but some adverse effects including mild neurological (e.g., headache, drowsiness, dizziness) and gastrointestinal symptoms (e.g., nausea, indigestion) were increased. In another systematic review study, Proctor et al. [20] reported that COX-2 specific inhibitors effectively reduced dysmenorrhea, but they are not used in many countries due to cardiovascular effects.

Potential side effects of some drugs make some women with dysmenorrhea to seek alternative medicine such as herbal drugs, diet, fish oil supplements, vitamin E, and low-fat diets and vegetables to control their symptoms [21–23]. Two Cochran systematic reviews investigated the efficacy of dietary supplements including vitamin and herbal drugs on dysmenorrheal pain relief. They found that primary studies had low quality evidence to support the effectiveness of any dietary supplement for dysmenorrhea, and there was a lack of evidence of their safety [24, 25].

Besides the above-mentioned relative risk factors of dysmenorrhea, the potential role of women's daily diet in association with the severity of dysmenorrhea has been reported. Abdul-Razzak et al. [21] reported the positive effect of calcium on controlling primary dysmenorrhea. Di Cintio et al. [26] showed a poor positive relationship between egg and cheese consumption and dysmenorrhea. Balbi et al. [27] believed that an excessive consumption of fish, eggs, and fruits had a negative relationship with the reduction of dysmenorrhea. A study at Dumlupinar University reported that coffee consumption and family history of dysmenorrhea were some significant risk factors of dysmenorrhea [28]. In the study by Ozerdogan et al.

[29], the prevalence of dysmenorrhea was higher in students who consumed tea, Coca cola, or Pepsi. The severity of dysmenorrhea was higher in women with excessive sugar intake.

Given the probable relationship between the nutritional status (excluding supplements) and primary dysmenorrhea, as well as significance of knowledge of nutrition-related methods for the management of dysmenorrhea, this systematic review investigated the relationship between dietary habits and nutritional status with the intensity of primary dysmenorrhea.

Methods

Search Process

In accordance with the PRISMA guideline for designing and implementing systematic review studies [30], the following steps were taken to perform this study: systematic literature search, organization of documents for review, abstracting and quality assessment of documents, synthesizing data, and writing the final report.

Search Strategy

A systematic literature search was performed in databases including Web of Science, Embase, Scopus, and PubMed (including Medline). The Boolean search method was used using keywords that were connected together via AND, OR, and NOT: ([Risk factor OR Associate factor OR contributing factor] OR [Food OR nutrient OR Diet OR Nutrition Status OR Nutrition Assessments OR Nutrition OR Nutrition Indexes OR Nutritional Indices OR Prognostic Nutritional Index OR Mini Nutritional Assessment]) AND (Dysmenorrhea OR Menstruating Disturbances OR Menstrual Disorders OR Menstrual Disorder OR Pelvic Pain OR Painful Menstruation OR Painful Period OR Period Pain OR Primary Dysmenorrhea OR Menstrual Pain OR Menstrual Cramps). Also, the reference list of included studies and previous systematic reviews were searched to increase the search coverage.

Inclusion Criteria

Type of Studies

Observational studies including cross-sectional, case-control, and cohort studies published in English language from 1990 to April 2018 were included. The studies published before 1990 and in languages other than English were excluded. Moreover, case reports, letters to the editor, and review articles were excluded. Trial studies on dietary supplements and herbal drugs were excluded because of the presence of a recent published systematic review on this topic [24].

Subjects

They were from all age ranges which were suffering from primary dysmenorrhea in their menstrual cycles. Those subjects with secondary dysmenorrhea who used contraceptive methods such as intrauterine contraceptive device or oral contraceptive pills were excluded. If the type of dysmenorrhea was not specified (due to a higher prevalence of primary dysmenorrhea in comparison to secondary dysmenorrhea), data was included.

Exposure

Observational studies on nutritional habits, daily food consumption, or nutrients in the daily diet were included. Trial studies on dietary supplements and herbal drugs were excluded.

Outcomes Measure

Intensity of dysmenorrheal pain was assessed using a visual analogue scale. Also, other valid scales or dichotomous measurements were other main outcomes measured in this systematic review.

Study Selection

The title and abstract of all articles retrieved during the search process were studied using the inclusion criteria. The full texts of the articles were examined using the above-mentioned criteria independently by 2 authors. Disagreements were resolved through discussions.

Quality Assessment

The modified STROBE checklist was used to assess the quality of selected observational studies with cohort, case-control, and cross-sectional designs. The STROBE as a valid instrument with 22 items was used for reporting or evaluating observational studies [31]. For the quality assessment of this review, the modified STROBE checklist with 10 items was used in relation of the methodological assessment of studies. The modified checklist consisted of objectives (specific objectives), study design (key elements of the study design), setting (the setting, location, relevant date, period of recruitment, exposure, follow-up, and data collection), subjects (eligibility criteria and the sources and methods of samples' selection), definition and measurement of all variables (outcomes, exposures, predictors for each variable of interest, sources of data, and methods of measurement), potential sources of bias and confounders (efforts taken to address potential sources of bias), sample size (final sample size), quantitative variables (handling variables during analyses, group selections, and reasons), and statistical methods (methods used to control confounding variables, methods used to examine subgroups and interactions, addressing missing data). Next, the studies were scored as "low risk", if the above mentioned criteria were unmet they were scored as "high risk", and if these items were not explained at all, they were rated as "not clearly defined". The quality assessment process was performed independently by 2 authors and disagreements were resolved through discussions.

Data Extraction

After screening and examining the quality of selected studies, data was extracted and recorded in forms independently by 2 authors. The forms were consisted of questions about the name of the first author, year, study design, country of the study, sample size, age of subjects, associations between dysmenorrhea and nutritional groups, and nutritional habits.

Data Synthesis

The qualitative synthesis of findings was performed due to the heterogeneity of measures used to assess and report nutrition consumption and insufficiency in provided data. Also, different measures were used to report the intensity of dysmenorrhea.

Findings

The search process led to retrieving 5,814 potentially relevant articles. Of which, 157 articles were found related to the study topic during abstract and title checking, but 18 articles were duplicated and 3 articles were published in other languages than English. The full texts of 136 remaining articles were reviewed and 38 articles were selected based on the inclusion criteria. The reference list of the included studies was read to find potentially related studies, but no article was included in the search process. Figure 1 showed the search process according to the PRISMA flowchart.

Description of Selected Studies

Most articles ($n = 27$) had cross-sectional designs, 4 had prospective or cohort designs, and the remaining were case-control studies. A total of 19,626 women were studied in these studies. The largest number of samples was 2,166 individuals in the study by Fujiwara and Nakata [32] in Japan. Also, the study of Abdul-Razzak et al. [33] in Jordan had the lowest number of samples ($n = 56$). The studies were performed in the different parts of the world including the United States, China, Japan, Iran, Jordan, Nepal, Turkey, India, South Korea, Ethiopia, Georgia, Serbia, and Italy. Japan had the largest number of studies with 6 articles (Table 1).

Quality Assessment

Evaluation of the selected studies using the modified checklist of STORBE with 10 items showed that the studies mostly had a moderate quality level. The quality assessment scores were presented in Table 1. The majority of the studies had issues in reporting bias, sample size estimation, description of design, and statistical analysis of quantitative variables. Figure 2 showed the graphical representation of the quality of studies.

Outcome Measurement

The findings of the selected studies were presented in 2 parts: (i) summarizing the results of the studies regarding the association between dysmenorrhea and dietary nutritional groups based on the food pyramid (Table 2) and (ii) abstracting the results of the studies based on nutritional and dietary habits (Table 3).

The Results of the Selected Studies According to the Dietary Nutritional Group

Carbohydrate Including Bread and Cereals: Consumption of pasta, rice, white bread, biscuits, pastry, potatoes, grains were studied by Di Cintio et al. [26], Balbi et al. [27], Tavallae et al. [16], and Gagua et al. [6], but no association was reported between the consumption of food related to these groups and dysmenorrhea.

Fruits and Vegetables: The association between the consumption of fruits and vegetables were investigated in 4 studies, but Di Cintio et al. [26] and Abu Helwa et al. [55] found no associations. However, Tavallae et al. [16] and Balbi et al. [27] reported that fruit consumption had negative associations with the level of menstrual pain.

Meat, Legumes, Eggs, and Nuts: This group consisted of a wide range of nutrients including meat, poultry, fish, legumes such as beans and soybean, eggs, and nuts. Fish consumption as a separate nutrient was studied by Deutch [34], Balbi et al. [27], and Grandi et al. [44] which reported inconsistent results. Deutch [34] found that a high intake of n-3 fatty acids through marine sources was




Fig. 1. PRISMA flow diagram.

associated with pain reduction; also Balbi et al. [27] found that women with dysmenorrhea significantly consumed less fish compared to healthy controls, but Grandi et al. [44] did not find any significant association. Egg as another nutrient was studied by Di Cintio et al. [26] and Balbi et al. [27] with completely contradicting results. Di Cintio et al. [26] reported that women with dysmenorrhea consumed eggs more frequently than healthy controls, but Balbi et al. [27] reported that women with dysmenorrhea consumed significantly less eggs compared to healthy controls. Other studies did not find significant results concerning the association between dysmenorrhea and the consumption of meat and its alternatives [16, 26, 27, 39, 52, 55].

Milk and Dairy: Milk and dairy consumption were reported in several studies. Four studies found a significant inverse association between the consumption of dairy products and dysmenorrhea [16, 21, 33, 59], Di Cintio et al. [26] found no relationship between the intake of milk and risk of dysmenorrhea, but cheese was more frequently used by women with dysmenorrhea than healthy controls.

Other Products: Other products included sugar, fat, salt, caffeine, vitamins, and minerals:

A) **Sugar:** Consumption of sugar in 3 studies had a significant relationship with dysmenorrhea. It developed a significantly higher risk of dysmenorrhea in women consuming excessive amount of sugar [6, 29, 60]. Inconsistently, Tavallae et al. [16], Grandi et al. [44], Mohamadirizi and Kordi [52], Abu Helwa et al. [55] found no association between the intake of sugar and dysmenorrhea.

B) **Fats:** Fat consumption was reported by 2 studies. Nagata et al. [39] found that only the intake of saturated fat had a positive association with lower menstrual pain. Also, Tavallae et al. [16]

reported that women with a low fat diet significantly experienced less pain during menstrual periods.

C) **Salt:** Three studies examined the relationship between excessive salt intake and dysmenorrhea. One study reported a higher prevalence of dysmenorrhea in women with moderate or excessive intake of salt compared with those with a minimal intake [29]. Two other studies did not report a significant association between a high salt intake and dysmenorrhea [44, 55].

D) **Caffeine Consumption:** Caffeine consumption was the most studied nutrient including coffee, tea, Nescafé, carbonated soft drinks, chocolate, and hot chocolate. In 15 studies, the association between caffeine intake and dysmenorrhea was assessed. In some studies, some forms of caffeinated drinks were assessed separately such as tea, coffee, Nescafé, but others studies assessed this group as whole. Seven studies [16, 29, 38, 44, 48, 52, 55] did not find any significant association between caffeine intake and dysmenorrhea, but Faramarzi and Salmalian [49], Pejčić and Jankovic [57], and Hailemeskel et al. [56] found significant associations between caffeine consumption and higher intensity of menstrual pain. Drinking tea did not have a significant association with dysmenorrhea in 4 studies [28, 43, 45, 60], but Wang et al. [37] reported more tea consumption among women with dysmenorrhea. The association between coffee and menstrual pain was reported in 3 studies [28, 43, 60] indicating a positive association, but 7 studies [45] reported that coffee consumption had no association with dysmenorrhea.

Vitamins and Minerals: Venkata et al. [42] reported reduced blood levels of Vitamin E and C and primary dysmenorrhea. Consistently, Pramanik et al. [53] reported a higher daily intake of beta carotene, vitamin-E, and zinc in girls without dysmenorrhea than

Table 1. Characteristics of selected primary studies

No.	Author, year	Study Design	Country	No. of participants	Age group, years	STROBE score	Outcome measured	Nutritional assessment
1	Deutch [34], 1995	Cross sectional	Denmark	181	20–45	6	Menstrual pain: have/not have	Two (prospective) 4-day dietary records were used to estimate average daily nutrient intake
2	Montero [35], 1996	Cross sectional	Spain	1147	14–20	8	Self-report of perceived pain as severe/medium/mild/no pain	Asked whether they had tried to lose weight
3	Di Cintio [26], 1997	Case control	Italy	251	Median: 26–27	8.5	Andersch and Milsom's classification on dysmenorrhea	Weekly frequencies of intake of each food item without information on portion size
4	Balbi et al. [27], 2000	Cross sectional	Naples	356	14–21	8	Intensity of the pain assessed with VAS	Frequency which certain foodstuffs were consumed
5	Fujiwara [36], 2003	Case control	Japan	439	18–20	7.5	Intensity of dysmenorrhea classified into score 1: free of pain or painful, but without need for analgesic; score 2: painful, requiring analgesic; score 3: painful, not relieved by analgesic	Grouped as having breakfast every morning, having breakfast 1–6 times per week, having breakfast less than once a week)
6	Wang et al. [37], 2004	Cohort	China	388	20–34	10	Menstrual pain (yes or no)	Tea consumption but not mentioned how measure
7	Weissman et al. [17], 2004	Cohort	USA	404	19–46	8	Four-point scale: no pain, minimal (can work somewhat uncomfortable), moderate (can work, but quite uncomfortable), or severe (miss work, have to be in bed)	Ask about vitamins use to cope with dysmenorrhea
8	Chung et al. [38], 2005	Cross sectional	Taiwan	151	21–44	7	Experience of dysmenorrhea as no/seldom/always	Drinking coffee habit as yes/no/quit
9	Nagata et al. [39], 2005	Cross sectional	Japan	276	19–24	8	Severity of menstrual pain using Andersch and Milson scoring system	Intakes of soy, fat and dietary fiber by assessing average frequency consumption of 169 food during the year
10	Fujiwara et al. [40], 2007	Cross sectional	Japan	716	18–20	6.5	Intensity of dysmenorrhea classified into score 1: free of pain or painful, but without need for analgesic; score 2: painful, requiring analgesic; score 3: painful, not relieved by analgesic	Dietary habits were classified into having no experience with dieting/currently on a diet/with a history of dieting
11	Ozerdogan et al. [29], 2009	Cross sectional	Turkey	857	17–32	8	Severity of menstrual pain using Andersch and Milson scoring system	Sugar and salt consumption categorized into minimal, moderate, or excessive daily intake. And caffeine consumption as infrequently/frequently

Table 1 (continued)

No.	Author, year	Study Design	Country	No. of participants	Age group, years	STROBE score	Outcome measured	Nutritional assessment
12	Fujiwara et al. [40], 2009	Cross sectional	Japan	315	18–20	7	Intensity of dysmenorrhea classified into score 1: free of pain or painful, but without need for analgesic; score 2: painful, requiring analgesic; score 3: painful, not relieved by analgesic	Categorized in 6 groups: I/ eating breakfast; II/skipping breakfast; III/not eating fast foods; IV/eating fast foods; V/not eating processed foods; VI/eating processed foods
13	Fujiwara and Nakata [41], 2010	Prospective	Japan	945	18–20	7.5	Intensity of dysmenorrhea classified into score 1: free of pain or painful, but without need for analgesic; score 2: painful, requiring analgesic; score 3: painful, not relieved by analgesic	Group I: Having breakfast every morning; Group II: having breakfast 1–6 times a week; and group III, having breakfast less than once a week
14	Abdul-Razzak et al. [21], 2010	Cross sectional	Jordan	127	19–24	7.5	Pain severity as mild: pain that resolved without the need for medication. Severe: pain that is resolved with simple analgesics Very severe: pain that is not relieved with simple analgesics and may interfere with usual daily activities	Frequency and type of dairy product intake (milk, yogurt, cheese, and labanah) recorded based on daily intake of dairy products
15	Unsal et al. [28], 2010	Cross sectional	Turkey	623	17–30	8	Severity of dysmenorrhea assessed by VAS and Andersch and Milson scoring system	Tea consuming: at least 4 glasses of tea in a day (75 mL), coffee consuming: at least 3 cups of coffee in a day (150 mL), cola consuming: at least a glass of cola in a day (200 mL), chocolate eating: at least 2 bars of chocolate in a day (150 mL)
16	Venkata et al. [42], 2011	Case control	China	191	18–23	4.5	Having dysmenorrhea: yes/no	Plasma vitamin C, plasma vitamin E
17	Bin Mahmud et al. [43], 2014	Cross sectional	Saudi Arabia	384	17–45	9.5	Pain as one of menstrual symptoms	Asked about usual daily consumption of the caffeinated beverages coffee, tea, Nescafe, carbonated soft drinks, chocolate and hot chocolate, in number of cups or cans per day
18	Tavallaee et al. [16], 2011	Cross sectional	Iran	381	16–56	7.5	Severity of dysmenorrhea assessed by Anderch and Milson scoring system	Consumption of fruits and vegetables/milk products/ meat and alternatives/ grains/fats and oils/ sweets and chocolates/ caffeine but not mentioned how measured
19	Gagua et al. [6], 2012	Cross sectional	Georgia	431	14–20	7	Having dysmenorrhea: yes/no	Asked about meal skipping: yes/no
20	Grandi et al. [44], 2012	Cross sectional	Italy	408	Mean 22.9	9.5	Pain intensity using a VAS	Sugar and salt consumption (minimal, moderate, excessive) and breakfast frequency (daily, 2/3 times a week, never)
21	Seven et al. [45], 2013	Case control	Turkey	380	18–23	7.5	Pain intensity using a VAS	Eating chocolate: yes /no

Table 1 (continued)

No.	Author, year	Study Design	Country	No. of participants	Age group, years	STROBE score	Outcome measured	Nutritional assessment
22	Fujiwara and Nakata [32], 2013	Prospective	Japan	2,166	18–20	5.5	Intensity of dysmenorrhea classified into score 1: free of pain or painful, but without need for analgesic; score 2: painful, requiring analgesic; score 3: painful, not relieved by analgesic	Skipping breakfast: yes/no
23	Vani et al. [46], 2013	Cross sectional	India	853	13–19	8	Occurrence of dysmenorrhea and if present, is it severe enough to skip classes or any need to take medications	Junk food consumption was assessed according frequency of eating junk per week and Fussy behavior at eating food and eating less food in order to lose weight
24	Gangwar et al. [47], 2014	Cross sectional	India	101	18–25	6	Pain intensity using a VAS	Diet as vegetarian/non-vegetarian
25	Sahin et al. [48], 2014	Cross sectional	Turkey	520	17–25	5.5	Pain intensity using a VAS	Consumption of tea/coffee/cola/chocolate as yes/no
26	Abdul-Razzak et al. [33], 2014	Cross sectional	Jordan	56	17–24	6	Pain intensity using a VAS and graded as mild/moderate/ severe/very sever	Frequency and type of dairy product intake (milk, yogurt, cheese, and labanah) recorded on daily basis
27	Faramarzi and Salmalian [49], 2014	Cross sectional	Iran	360	17–25	8	Graded as 0 = absence of dysmenorrhea. 1 = mild pain, 2 = moderate pain, 3 = severe pain	Caffeine intake defined as excessive if consumption of caffeinated soft drinks, coffee, decaffeinated coffee, tea, chocolate milk, and chocolate bars in the daily diet was self-reported as ≥300 mg/day and otherwise it was as minimum to moderate. Having breakfast was classified as normal if participant had breakfast one to six times per week and as low if she had breakfast less than once a week
28	Jeon et al. [50], 2014	Cross sectional	South Korea	572	11–16	8.5	Perception of menstruation as discomfort/painfulness/none	Ask about regular/irregular eating
29	Kazama et al. [51], 2015	Cross sectional	Japan	1,018	12–15	6.5	Pain intensity using a VAS	Habitual breakfast, lunch, and supper classified as (1) everyday, (2) sometimes, and (3) none
30	Mohamadirizi and Kordi [52], 2015	Cross sectional	Iran	407	10–15	9	Pain as one dimension of menstrual distress questionnaire	The nutrition pattern questionnaire included 4 main nutrients: sweet-fatty (chocolates, cake, ice cream, nuts, etc.), salty-fatty (soya, red meat, chicken meat, etc.), fast foods (hamburger, pizza, potato chips, etc.), and caffeine-containing materials (tea, coffee, etc.). The frequency of their consumption within the past year was investigated

Table 1 (continued)

No.	Author, year	Study Design	Country	No. of participants	Age group, years	STROBE score	Outcome measured	Nutritional assessment
31	Pramanik et al. [53], 2015	Case control	India	379	12–17	6.5	Having dysmenorrhea: yes/no	24 h food consumption recall
32	Bavil et al. [54], 2016	Cross sectional	Iran	250	18–26	7.5	Having dysmenorrhea: yes/no	Eating behavior but not clearly explained how measured
33	Abu Helwa et al. [55], 2018	Cross sectional	Palestine	956	Mean 19.7	8.5	Pain intensity using a VAS	Nutritional habits as having breakfast every day/ sometimes/never and habits about consumption of tea/ coffee/nescafe (≤ 1 cup or ≥ 2 cup)/ consumption of Vegetables and fruits, meat and protein rich diet, sweets and sugary beverages, salty foods graded as never or low/moderate/high/very high over last 6 months
34	Hailemeskel et al. [56], 2016	Cross sectional	Ethiopia	440	Mean 20.6	7.5	Severity of dysmenorrhea assessed by VAS and Anderch and Milson scoring system	Ask about history of attempt to lose weight: yes/no
35	Pejčić and Jankovic [57], 2016	Case control	Serbia	288	18–29	9.5	Dysmenorrhea yes/no	Ask about Consuming at least one cup of coffee daily
36	Shinde and Laddad [58], 2016	Cross sectional	India	624	11–19	6	Dysmenorrhea yes/no	Frequency of eating junk food (categorized into <3 days per week and >3 days per week
37	Karacin et al. [59], 2018	Case control	Turkey	683	18–25	9	Severity of dysmenorrhea assessed by VAS	The intake of dairy products assessed on daily basis
38	Muluneh et al. [60], 2018	Cross sectional	Ethiopia	539	14–24	9.5	Dysmenorrhea yes/no	Sugar intake: Excessive if individuals took 12 or more teaspoons of table sugar daily, Moderate if 6 to 12 teaspoons; and in a restricted use if less than 6 teaspoons




Fig. 2. Quality assessment of selected studies according to STROBE checklist.

Table 2. Summarized results of reviewed studies according to dietary nutritional group and habit

Dietary nutritional group	Related studies	Nutritional assessment	Relation with dysmenorrhea
Carbohydrate including bread and cereals	Di Cintio et al. [26], 1997	Portions of pasta, rice, white bread, biscuits, pastry, potatoes per week	No relationship with risk of dysmenorrhea.
	Balbi et al. [27], 2000	Consumption of pasta	No significant difference between cases and healthy controls
	Tavallae et al. [16], 2011	Consumption of grains	No association with menstrual pain level
	Gagua [6], 2012	Excessive consumption of bread	No association with dysmenorrhea
Fruits and vegetables	Di Cintio et al. [26], 1997	Portions of different kinds of vegetables, green vegetables and fresh fruits per week	No relationship with risk of dysmenorrhea.
	Balbi et al. 2000 [27]	Consumption of fruit	Significantly lower fruit consumption in dysmenorrheic cases compared to healthy control
	Tavallae et al. [16], 2011	Consumption of fruits and vegetables	Significant inverse association between pain level and fruit and vegetable consumption
	Abu Helwa [55], 2018	Consumption of fruits and vegetables	No significant difference
Meat, legumes, and nuts	Di Cintio et al. [26], 1997	Portions of beef, pork, poultry, other meats (rabbit, lamb), fish, liver, raw and boiled ham, salami, sausage, tinned meat per week	No relationship between intake of these dietary factors and risk dysmenorrhea
	Balbi et al. [27] 2000	Consumption of meat	No difference in meat consumption in case and controls
	Nagata et al. [39], 2005	Soy products in daily dietary	No significant association with dysmenorrhea
	Tavallae [16], 2011	Consumption of meat and alternatives	No association with menstrual pain level
	Mohamadirizi and Kordi [52], 2015	Consumption of salty-fatty foods including soya, red meat, chicken meat, etc	No association with dysmenorrhea
	Abu Helwa [55], 2018	Meat and protein rich diet	No significant difference according to dysmenorrhea
Fish	Deutch [34], 1995	N-3 fatty acid intake from marine source	Decrease in pain with a high n-3 fatty acid intake
	Balbi et al, [27]2000	Fish consumption	Significantly lower fish consumption in dysmenorrheic cases compared to healthy control
	Grandi et al. [44], 2012	Fish consumption	No association with dysmenorrhea
Eggs	Di Cintio et al. [26] 1997	Portions of eggs per week	Eggs were more frequently used by dysmenorrheic cases than healthy controls
	Balbi et al, [27], 2000	Egg consumption	Significantly lower eggs consumption in dysmenorrheic cases compared to healthy control
Milk and dairy	Di Cintio et al. [26], 1997	Portions of milk and cheese per week	No relationship between intake of milk and risk dysmenorrhea but cheese was more frequently used by cases than controls
	Abdul-Razzak et al, [21], 2010	Dietary intake of dairy products as frequency and type of diary product intake including milk, yogurt, cheese, and labanah on a daily basis as none, 1, 2, 3, or up to 4 dairy servings per day	Increased intake of dairy of three to 4 servings a day was significantly associated with reduction in dysmenorrhea
	Tavallae [16], 2011	Consumption of milk products	Significant inverse association with menstrual pain

Table 2 (continued)

Dietary nutritional group	Related studies	Nutritional assessment	Relation with dysmenorrhea	
	Abdul-Razzak et al, [33], 2014	Frequency and type of dairy product intake (milk, yogurt, cheese, and labanah)	There was significant reduction in prevalence of very severe menstrual pain by increasing daily dairy servings.	
	Karacin et al. [59], 2018	Consumption of dairy products	The dysmenorrhea group had significantly less consumption of dairy products	
Other products	Sugar	Ozerdogan et al. [29], 2009	Sugar intake based on self-reported excessive consumption of bread, table sugar, cola, chocolate, rice, carrots, ice cream, cake, yoghurt, noodles, milk, and sweets in daily diet	Significantly higher risk of dysmenorrhea in women consuming excessive sugar compared with women with low intake
		Tavallaee et al. [16], 2011	Consumption of sweets and chocolate	No association with menstrual pain was found
		Gagua et al. [6], 2012	Excessive sugar intake based on individual self-report of excessive consumption of table sugar, soda, chocolate, ice cream, and sweets in their daily diet	Increased dysmenorrhea was associated with increased intake of sugar
		Grandi et al. [44], 2012	Excessive sugar consumption	No association with dysmenorrhea
		Mohamadirizi and Kordi [52], 2015	Consumption of sweet-fatty foods (chocolates, cake, ice cream, nuts, etc.)	No association with dysmenorrhea
		Abu Helwa [55], 2018	Sweets and sugary beverages	No significant difference according to dysmenorrhea
		Muluneh et al. [60], 2018	Sugar intake	Statistically increased risk of dysmenorrhea was associated with sugar intake
	Fat	Nagata [39], 2005	Total fat, saturated fat, monounsaturated fat, polyunsaturated fat	Only saturated fat intake had positive association with lower menstrual pain
		Tavallaee, [16] 2011	Consumption of fat and oil	Women who tended to have a low fat diet significantly experienced less pain during their menstrual periods
	Salt	Ozerdogan [29], 2009	Excessive salt intake	Significantly higher prevalence of dysmenorrhea in women who reported consuming a moderate or excessive intake of salt compared with women who had a minimal intake
		Grandi et al. [44], 2012	Excessive salt consumption	No association with dysmenorrhea
		Abu Helwa, [55], 2016	Salty foods	No significant difference according to dysmenorrhea
	Caffeine	Wang [37], 2004	Tea consumption	Significantly more tea consumption among dysmenorrheic women
Chung [38], 2005		Coffee consumption	No correlation between coffee consumption and dysmenorrhea.	
Ozerdogan [29], 2009		Intake of caffeine including caffeinated soft drinks, coffee, decaffeinated coffee, tea, chocolate milk, and chocolate bars	No correlation between caffeine consumption and dysmenorrhea.	
Unsal et al. [28], 2010		Consumption of coffee, tea, cola, and chocolate	Significant correlation of dysmenorrhea with coffee consumption but not tea, cola and Chocolate consumption	
Bin Mahmoud [43], 2011		Daily consumption of the caffeinated beverages including coffee, tea, Nescafe, carbonated soft drinks, chocolate and hot chocolate, in number of cups or cans per day	Nescafe was a risk factor for menstrual symptoms but No significant risk seen with drinking tea or carbonated soft drinks	

Table 2 (continued)

Dietary nutritional group	Related studies	Nutritional assessment	Relation with dysmenorrhea
	Tavallae et al. [16], 2011	Coffee consumption	No association with dysmenorrhea
	Grandi et al. [44], 2012	Coffee consumption	No association with dysmenorrhea
	Seven [45], 2013	Coffee, tea and chocolate consumption	Coffee and tea had no association with dysmenorrhea, but chocolate eating significantly affect dysmenorrhea
	Sahin [48], 2014	Cola, tea, coffee, chocolate consumption	No association with dysmenorrhea
	Faramarzi [49], 2014	Caffeine intake including caffeinated soft drinks, coffee, decaffeinated coffee, tea, chocolate milk, and chocolate bars in the daily diet	High caffeine intake had significant association with dysmenorrhea
	Mohamadirizi and Kordi [52], 2015	Consumption of caffeine-containing materials (tea, coffee, etc.).	No association with dysmenorrhea
	Abu Helwa [55], 2018	Tea/coffee/Nescafe	No association with dysmenorrhea
	Pejčić and Jankovic [57], 2016	Daily coffee intake	Significantly associated with dysmenorrhea
	Hailemeskel et al. [56], 2016	Tea, Coca-Cola or Pepsi and Chocolate consumption per day	Significantly associated with dysmenorrhea
	Muluneh et al. [60], 2018	Daily coffee and tea intake	Tea was not significantly associated but coffee more than 5 cups per day was significantly associated with dysmenorrhea
Vitamins and minerals	Weissman et al. [17], 2004	Vitamin use	No significant association between vitamin use and intensity of dysmenorrhea
	Venkata [42], 2011	Vitamin E & C blood level	Significantly decreased blood levels of Vitamin E, C in cases with primary dysmenorrhea
	Abdul-Razzak et al. [33], 2014	Plasma vitamin D concentration	A high prevalence of vitamin D insufficiency in participants with severe and very severe dysmenorrhea.
	Pramanik et al. [53], 2015	Daily intake of beta carotene, vitamin-E, vitamin-C and zinc	Significantly higher daily intake of beta carotene, vitamin-E and zinc in girls without dysmenorrhea than dysmenorrheic counterpart. But no difference according to vitamin C
Fiber	Nagata et al. [39], 2005	Dietary fiber	Negative and significant association between dietary fiber and intensity of dysmenorrhea

their counterparts with dysmenorrhea. No difference was reported with vitamin C. Weissman et al. [17] reported a lack of significant relationship between vitamin intake and dysmenorrhea. Only Abdul-Razzak et al. [33] reported that individuals with severe and very severe dysmenorrhea had vitamin D deficiency or had low calcium intake. Nevertheless, there was no relationship between the severity of menstrual pain, and vitamin D and plasma calcium levels.

Fiber Consumption: A study examined the association between fiber consumption and dysmenorrhea, and reported a negative and significant relationship between them [39].

Results of the Studies According to Nutritional Habits Following Diet to Lose Weight: Montero et al. [35], Fujiwara [61], Rupa Vani et al. [46], and Hailemeskel et al. [56] studied the association between the current or previous attempts of going on a diet to lose weight and the intensity of dysmenorrhea. Consistently, a positive association between following a diet to lose weight and dysmenorrhea was reported.

Following Vegetarian Diet: Only one study focused on the relationship between having a vegetarian diet and dysmenorrhea. No significant difference was found in the prevalence of dysmenorrhea in vegetarian and non-vegetarian subjects [47].

Table 3. Summarized results of reviewed studies according to nutritional habits

Nutritional habit	Related studies	Nutritional assessment	Relation with dysmenorrhea
Following diet to lose weight	Montero et al. [35], 1996	Dieting for loose weight	Attempting to lose weight was significantly associated with increased menstrual pain
	Fujiwara [61], 2007	Having no experience with dieting/ currently on a diet/having history of dieting	Participants with history of dieting had higher intensity of dysmenorrhea than the other groups
	Rupa Vani et al. [46], 2013	Dieting for lose weight	Significantly associated with dysmenorrhea
	Hailemeskel et al. [56], 2016	History of attempt to lose weight	Significantly associated with dysmenorrhea
Following vegetarian diet	Gangwar et al. [47], 2014	Dietary habits as vegetarian/ non-vegetarian	No significant difference in the prevalence of dysmenorrhea in vegetarian and non-vegetarian subjects
Eating behavior	Fujiwara [36], 2003	Skipping breakfast	A significant correlation between skipping breakfast and intensity of dysmenorrhea
	Fujiwara et al. [40], 2009	Eating/skipping breakfast; eating/not eating fast foods eating/ not eating processed foods	Dysmenorrhea scores were significantly higher in participants skipping breakfast, who ate fast foods and processed foods compared to their counterparts.
	Fujiwara and Nakata [41], 2010	Having breakfast every morning/having breakfast 1–6 times a week/having breakfast less than once a week	Severity of dysmenorrhea was significantly higher in the population that skipped breakfast
	Gagua et al. [6], 2012	Meal skipping based on number of meal intakes throughout the 24 h	Meal skipping (inadequate nutrition) was one of most important risk factors of dysmenorrhea
	Grandi et al. [44], 2012	Attitude to eating breakfast	No association with dysmenorrhea
	Fujiwara and Nakata [32], 2013	Skipping breakfast	Significantly higher severity of dysmenorrhea in women that skipped breakfast
	Rupa Vani et al. [46], 2013	Eating junk foods	No association with dysmenorrhea
	Faramarzi and Salmalian [49], 2014	Having breakfast	No significant relation with dysmenorrhea
	Jeon et al. [50], 2014	Regular/Irregular eating	The score for dysmenorrhea was significantly higher when the participants ate meals irregularly
	Kazama et al. [51], 2015	Having breakfast	Likelihood of experiencing dysmenorrhea significantly increased when sometimes eating breakfast or skipped breakfast
	Mohamadirizi and Kordi [52], 2015	Having fast foods like hamburger, pizza, potato chips, etc.	No significant relation with dysmenorrhea
	Bavil et al. [54], 2016	Eating behavior	Significant differences between the groups with and without dysmenorrhea in terms of eating behavior
	Abu Helwa [55], 2018	Skipping breakfast	Skipping breakfast was the only significant predictors of moderate/severe dysmenorrhea
Shinde and Laddad [58], 2016	Eating junk foods	Dysmenorrhea were more frequent in girls with habit of eating junk food <3 days a week	

Eating Behavior: Eating behaviors such as skipping breakfast, irregular eating, fast foods, or process foods were studied. Skipping breakfast was the most studied eating behavior and 6 studies found a significant correlation between skipping breakfast and the intensity of dysmenorrhea. Those women who skipped breakfast suf-

fered from a higher intensity of dysmenorrhea [32, 36, 40, 41, 51, 55]. Conversely, Faramarzi and Salmalian [49] did not find a significant relationship between eating breakfast and dysmenorrhea. Also, no association was observed between the attitude toward eating breakfast and dysmenorrhea [44].

Habits of regular eating or meal skipping were studied by Gagua et al. [6] and Jeon et al. [50]. They showed that those subjects with irregular and skipped meal habits had a significantly higher intensity of dysmenorrhea. Also Babil et al. [54] found significant differences between the groups with and without dysmenorrhea in terms of eating behaviors.

Studies on eating fast foods/process foods/junk foods reported inconsistent results. Fujiwara et al. [40] and Shinde and Laddad [58] reported a significantly higher level of dysmenorrhea in those subjects who ate fast foods and processed foods compared to their healthy counterparts. Mohamadirizi and Kordi [54] and Rupa Vani et al. [46] did not report such an association.

Discussion

Dysmenorrhea is one of the main factors influencing the quality of life and social activities of women [1, 8]. Several factors can affect primary dysmenorrhea and its incidence and severity. Nutrition has not been adequately studied with regard to its effect on dysmenorrhea [21, 27]. Therefore, this systematic review aimed to investigate the association between primary dysmenorrhea and dietary habits.

Regarding dietary habits and the consumption of different nutritional groups, the consumption of carbohydrates including bread, cereals, meat, legumes, and nuts had no association with dysmenorrhea and its intensity. Consumption of fish and eggs as the alternatives of meat showed inconsistent results. Regarding the association between fish and dysmenorrhea, the consumption of fish was associated with less dysmenorrhea pain, but 2 studies found inconsistent results. This finding is consistent with results of recent studies indicating the effects of fish oil on primary dysmenorrhea [62, 63]. The prevention of prostaglandins synthesis is the major mechanism of fish oil [20]. The protective role of fish is due to the content of omega-3 fatty acids. If more dietary omega-3 fatty acids are consumed, they enter the phospholipid membrane [64]. During menstruation, omega-3 fatty acids compete with fatty acids containing omega-6 to produce prostaglandins and leukotriene. In addition, they can hinder the production of arachidonic acid [27]. The effect of arachidonic acid in animal fats, especially meat, is the synthesis of prostaglandin [54]. Prostaglandins produced from omega-3 fatty acids are less potent, and can reduce the contraction of the myometrium and vessels [16]. A reduction of systemic symptoms is due to a poor leukotriene's production. Egg can prevent dysmenorrhea due to its magnesium and calcium contents. Magnesium such as calcium plays a key role in the membrane fixation mechanism and can modulate synapses through calcium-de-

pendent mechanisms. Reductions in magnesium levels increase synaptic transmission, resulting in myometric muscle contractions and, consequently, dysmenorrhea [27]. Foods containing the high levels of magnesium can reduce the severity of dysmenorrhea through reducing the synthesis of prostaglandins and spasticity of small muscles and vessels [54]. However, such findings are not consistent with that of the current systematic review. Such inconsistency may be due to small numbers of studies in this subgroup.

Studies on the association between the consumption of fruits and vegetables and dysmenorrhea were inconsistent. Two of 4 studies in this subgroup showed no significant association [26, 55], but 2 other studies confirmed that the increased consumption of fruits and vegetables are associated with the reduction of menstrual pain [16, 27]. Fruits and vegetables are important sources of vitamins and antioxidants [65]. Therefore, the combination of these results with the findings of studies on the plasma level of vitamins confirmed that women with dysmenorrhea took less fruits and vegetables and had less plasma levels of vitamins [33, 42, 53]. Vitamins, beta-carotene, zinc, and minerals are diet antioxidants. Vitamin E is fat soluble and is primarily responsible for oxidant membrane injuries and is the first line of defense against the peroxidation of phospholipids. Vitamin C (ascorbic acid) is soluble in water and provides an antioxidant capacity by removing oxygen-free radicals. So vitamin C can convert vitamin-E-free radical back to vitamin E and are essential for the recycling of vitamin E to prevent fat peroxidation. Beta carotene inhibits oxidants. Zinc is an antioxidant factor [53]. Reducing the level of antioxidants can increase the consumption of antioxidants or free radicals to detoxify the increased levels of oxidants in primary dysmenorrhea. A study on vitamins E and B (thiamine), magnesium, and fish oil suggests that these foods are important sources of vitamins E, A, and other nutrients [7]. Women with primary dysmenorrhea who use non-steroidal anti-inflammatory drugs are suggested to use natural antioxidants such as vitamin E, vitamin C, and β -carotenoids for the reduction of pain [7, 53]. Single randomized controlled trials showed the effect of vitamin supplements on the reduction of menstrual pain reduction [63, 66, 67], but a Cochrane meta-analysis on the effect of dietary supplements for primary dysmenorrhea lacked high quality evidence to support the effectiveness and safety of any dietary supplement for dysmenorrhea. Therefore, more research with higher methodological qualities are required [24].

Five studies assessed the association between the dairy consumption (calcium intake) and dysmenorrhea and

they mostly (4 out of 5) reported a positive association between dairy consumption and the reduction of menstrual pain [16, 21, 33, 59]. Calcium intake in the diet has a protective effect on dysmenorrhea [21]. Calcium is absorbed in the upper intestine and can regulate the ability of muscle cells to respond to nerve stimuli through various functions, which is called stabilizing [21, 27, 33]. Therefore, dairy and calcium consumption can reduce dysmenorrhea through controlling muscle nervous activities. Conversely, reduction of calcium concentrations can increase muscle spasm, spasticity, and muscle contractions [21]. The possible explanation for this relationship is that the absorption and metabolism of vitamins and minerals are endangered by the use of sugar leading to muscle spasm and menstrual pain [56].

Sugar consumption had an inconsistent association with dysmenorrhea. Excessive sugar intake was increased by the absorption of certain vitamins and minerals and led to a food instability. It creates muscle spasm and pain in the menstrual cycle. Also, certain nutrients directly influence the circulation status of sexual steroids and cause muscle spasm [6, 29, 48, 56]. The relationship between caffeine intake and dysmenorrhea has been reported in many studies due to its effect on vasoconstriction and pelvic pain [49]. On the other hand, caffeine can stimulate stress and irritability [28, 37, 49, 56].

Nutritional deficiencies are the most important factors that disrupt the hypothalamic-pituitary-ovarian axis. Nutritional deficiencies such as not eating breakfast contribute to the development of gynecological disorders especially endometriosis [36]. The nutritional status influences sexual steroids. Insufficient nutrition causes a change in hormonal levels due to reduced levels of energy [6]. Menstrual pain occurs only during ovulation. Progesterone affects the synthesis of prostaglandins and their attachment to myometrium receptors. Prostaglandins affect the uterine muscle and vascular tone, and dysmenorrhea is caused by an imbalance in prostaglandins. In addition, progesterone has some effects on the concentration of prostaglandin in myometrium [37].

The majority of the studies had issues in reporting potential source of bias, sample size estimation, description of design, and statistical analysis of quantitative variables. So there is a need to design and implement studies with stronger methodologies.

Limitations

The most important limitation of this review study was methodological heterogeneity of studies. The heterogeneity in the assessment and report of consumption of nu-

tritional groups and habits limited the data synthesis to qualitative synthesis. Performing meta-analysis was impossible. Qualitative synthesis has limitations in terms of providing conclusive findings. A lack of pre-registration protocol and exclusion of studies publications prior to 1990 and published in other languages are the other limitations of this study.

Conclusion

This systematic review showed the relationship between some nutritional factors and primary dysmenorrhea, indicating the need for further studies on this topic. The increased consumption of fruits and vegetables as the sources of many vitamins and minerals, fish and milk, and dairy can have a positive association with less menstrual pain. Other findings are related to nutritional habits, but most studies showed a negative association between dysmenorrhea and meal skipping and following diet to lose weight. Therefore, attention should be paid to the correction of female nutritional behaviors for having enough and balanced diet. Given the negative effects of dysmenorrhea on the quality of life, measures should be taken to increase the knowledge of women about dysmenorrhea and make appropriate lifestyle changes to control it. Designing and implementing interventional studies on nutrition education and modification are recommended.

Disclosure Statement

The authors declare no conflicts of interest.

Financial Support

None.

References

- 1 Iacovides S, Avidon I, Baker FC: What we know about primary dysmenorrhea today: a critical review. *Hum Reprod Update* 2015;21: 762–778.
- 2 Jonatan S, Berek J: *Novak's Gynecology: 14th*. New York, Lippincott Williams and Wilkins Publishers, 2012.
- 3 Chuamoor K, Kaewmanee K, Tanmahasamut P: Dysmenorrhea among Siriraj nurses; prevalence, quality of life, and knowledge of management. *J Med Assoc Thai* 2012;95: 983–991.

- 4 Mathias SD, Kuppermann M, Liberman RF, Lipschutz RC, Steege JF: Chronic pelvic pain: prevalence, health-related quality of life, and economic correlates. *Obstet Gynecol* 1996;87:321–327.
- 5 Shewte MK, Sirpurkar MS: Dysmenorrhoea and quality of life among medical and nursing students: a cross-sectional study. *Nat J Comm Med* 2016;7:474–479.
- 6 Gagua T, Tkeshelashvili B, Gagua D: Primary dysmenorrhea: prevalence in adolescent population of Tbilisi, Georgia and risk factors. *J Turk Ger Gynecol Assoc* 2012;13:162–168.
- 7 Haidari F, Akrami A, Sarhadi M, Mohammad Shahi M: Prevalence and severity of primary dysmenorrhea and its relation to anthropometric parameters. *Tums Hayat* 2011;17:70–77.
- 8 Ju H, Jones M, Mishra G: The prevalence and risk factors of dysmenorrhea. *Epidemiol Rev* 2014;36:104–113.
- 9 Burnett MA, Antao V, Black A, Feldman K, Grenville A, Lea R, et al: Prevalence of primary dysmenorrhea in Canada. *J Obstet Gynaecol Can* 2005;27:765–770.
- 10 Lathe P, Mignini L, Gray R, Hills R, Khan K: Factors predisposing women to chronic pelvic pain: systematic review. *BMJ* 2006;332:749–755.
- 11 Nohara M, Momoeda M, Kubota T, Nakabayashi M: Menstrual cycle and menstrual pain problems and related risk factors among Japanese female workers. *Ind Health* 2011;49:228–234.
- 12 Ohde S, Tokuda Y, Takahashi O, Yanai H, Hinohara S, Fukui T: Dysmenorrhea among Japanese women. *Int J Gynaecol Obstet* 2008;100:13–17.
- 13 Patel V, Tanksale V, Sahasrabhojane M, Gupte S, Nevrekar P: The burden and determinants of dysmenorrhoea: a population-based survey of 2262 women in Goa, India. *BJOG* 2006;113:453–463.
- 14 Pawłowski B: Prevalence of menstrual pain in relation to the reproductive life history of women from the Mayan rural community. *Ann Hum Biol* 2004;31:1–8.
- 15 Santer M, Warner P, Wyke S: A scottish postal survey suggested that the prevailing clinical preoccupation with heavy periods does not reflect the epidemiology of reported symptoms and problems. *J Clin Epidemiol* 2005;58:1206–1210.
- 16 Tavallaee M, Joffres MR, Corber SJ, Bayanzadeh M, Rad MM: The prevalence of menstrual pain and associated risk factors among Iranian women. *J Obstet Gynaecol Res* 2011;37:442–451.
- 17 Weissman AM, Hartz AJ, Hansen MD, Johnson SR: The natural history of primary dysmenorrhoea: a longitudinal study. *BJOG* 2004;111:345–352.
- 18 Osayande AS, Mehulic S: Diagnosis and initial management of dysmenorrhoea. *Am Fam Physician* 2014;89:341–346.
- 19 Marjoribanks J, Proctor M, Farquhar C, Derks RS: Nonsteroidal anti-inflammatory drugs for dysmenorrhoea. *Cochrane Database Syst Rev* 2015;7:CD001751.
- 20 Proctor M, Farquhar C: Diagnosis and management of dysmenorrhoea. *BMJ* 2006;332:1134–1138.
- 21 Abdul-Razzak KK, Ayoub NM, Abu-Taleb AA, Obeidat BA: Influence of dietary intake of dairy products on dysmenorrhoea. *J Obstet Gynaecol Res* 2010;36:377–383.
- 22 Proctor ML, Murphy PA: Herbal and dietary therapies for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2001;3:CD002124.
- 23 Sidani M, Campbell J: Gynecology: select topics. *Prim Care* 2002;29:297–321.
- 24 Pattanittum P, Kunyanone N, Brown J, Sangkomkamhang US, Barnes J, Seyfoddin V, et al: Dietary supplements for dysmenorrhoea. *Cochrane Database Syst Rev* 2016;3:CD002124.
- 25 Zhu X, Proctor M, Bensoussan A, Wu E, Smith CA: Chinese herbal medicine for primary dysmenorrhoea. *Cochrane Database Syst Rev* 2008;2:CD005288.
- 26 Di Cintio E, Parazzini F, Tozzi L, Luchini L, Mezzopane R, Marchini M, et al: Dietary habits, reproductive and menstrual factors and risk of dysmenorrhoea. *Eur J Epidemiol* 1997;13:925–930.
- 27 Balbi C, Musone R, Menditto A, Di Prisco L, Cassese E, D' Ajello M, et al: Influence of menstrual factors and dietary habits on menstrual pain in adolescence age. *Eur J Obstet Gynecol Reprod Biol* 2000;91:143–148.
- 28 Unsal A, Ayranci U, Tozun M, Arslan G, Calik E: Prevalence of dysmenorrhea and its effect on quality of life among a group of female university students. *Ups J Med Sci* 2010;115:138–145.
- 29 Ozerdogan N, Sayiner D, Ayranci U, Unsal A, Giray S: Prevalence and predictors of dysmenorrhea among students at a university in Turkey. *Int J Gynaecol Obstet* 2009;107:39–43.
- 30 Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- 31 von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP: The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med* 2007;4:e296.
- 32 Fujiwara T, Nakata R: Skipping breakfast may induce dysmenorrhoea and irregular menstruation in post-adolescent women in Japan. *Ann Nutr Metab* 2013;63:668.
- 33 Abdul-Razzak KK, Obeidat BA, Al-Farras MI, Dauod AS: Vitamin D and PTH status among adolescent and young females with severe dysmenorrhea. *J Pediatr Adolesc Gynecol* 2014;27:78–82.
- 34 Deutch B: Menstrual pain in Danish women correlated with low n-3 polyunsaturated fatty acid intake. *Eur J Clin Nutr* 1995;49:508–516.
- 35 Montero P, Bernis C, Fernandez V, Castro S: Influence of body mass index and slimming habits on menstrual pain and cycle irregularity. *J biosoc Sci* 1996;28:315–323.
- 36 Fujiwara T: Skipping breakfast is associated with dysmenorrhea in young women in Japan. *Int J Food Sci Nutr* 2003;54:505–509.
- 37 Wang L, Wang X, Wang W, Chen C, Ronnenberg AG, Guang W, et al: Stress and dysmenorrhoea: a population based prospective study. *Occup Environ Med* 2004;61:1021–1026.
- 38 Chung FF, Yao CC, Wan GH: The associations between menstrual function and life style/working conditions among nurses in Taiwan. *J Occup Health* 2005;47:149–156.
- 39 Nagata C, Hirokawa K, Shimizu N, Shimizu H: Associations of menstrual pain with intakes of soy, fat and dietary fiber in Japanese women. *Eur j clin nutr* 2005;59:88–92.
- 40 Fujiwara T, Sato N, Awaji H, Sakamoto H, Nakata R: Skipping breakfast adversely affects menstrual disorders in young college students. *Int J Food Sci Nutr* 2009;60:23–31.
- 41 Fujiwara T, Nakata R: Skipping breakfast is associated with reproductive dysfunction in post-adolescent female college students. *Appetite* 2010;55:714–717.
- 42 Venkata RS, Kiran VR, Vijayasree M: Oxidative stress and antioxidant status in primary dysmenorrhea. *J Clin Diagn Res* 2011 5:509–511.
- 43 Bin Mahmoud AZ, Makhdoom AN, Mufti LA, Alreheli RS, Farghal RG, Aljaouni SE: Association between menstrual disturbances and habitual use of caffeine. *J Taibah University Med Sci* 2014;9:341–344.
- 44 Grandi G, Ferrari S, Xholli A, Cannoletta M, Palma F, Romani C, et al: Prevalence of menstrual pain in young women: what is dysmenorrhea? *J Pain Res* 2012;5:169–174.
- 45 Seven M, Guvenc G, Akyuz A, Eski F: Evaluating dysmenorrhea in a sample of Turkish nursing students. *Pain Manag Nurs* 2013;15:664–671.
- 46 Vani R, Veena S, Subitha L, Hemanth Kumar R, Bupathy A: Menstrual abnormalities in school going girls – are they related to dietary and exercise pattern? *J Clin Diagn Res* 2013;7:2537–2540.
- 47 Gangwar V, Kumar D, Gangwar RS, Arya M, Banoo H: Prevalence of primary dysmenorrhea among the undergraduate medical students and its impact on their performance in study. *Int J Physiol* 2014;2:14.
- 48 Sahin S, Ozdemir K, Unsal A, Arslan R: Review of frequency of dysmenorrhoea and some associated factors and evaluation of the relationship between dysmenorrhoea and sleep quality in university students. *Gynecol Obstet Invest* 2014;78:179–185.
- 49 Faramarzi M, Salmalian H: Association of psychologic and nonpsychologic factors with primary dysmenorrhoea. *Iran Red Crescent Med J* 2014;16:e16307.

- 50 Jeon GE, Cha NH, Sok SR: Factors influencing the dysmenorrhea among Korean adolescents in middle school. *J Phys Ther Sci* 2014; 26:1337–1343.
- 51 Kazama M, Maruyama K, Nakamura K: Prevalence of dysmenorrhea and its correlating lifestyle factors in Japanese female junior high school students. *Tohoku J Exp Med* 2015;236: 107–113.
- 52 Mohamadirizi S, Kordi M: The relationship between food frequency and menstrual distress in high school females. *Iran J Nurs Midwifery Res* 2015;20:689–693.
- 53 Pramanik P, Banerjee SB, Saha P: Primary dysmenorrhea in school going adolescent girls – is it related to deficiency of antioxidant in diet? *Int J Life Sci Pharma Res* 2015;5:L54–L63.
- 54 Babil DA, Dolatian M, Mahmoodi Z, Baghban AA: Comparison of lifestyles of young women with and without primary dysmenorrhea. *Electron Physician* 2016;8:2107–2114.
- 55 Abu Helwa HA, Mitaeb AA, Al-Hamshri S, Sweileh WM: Prevalence of dysmenorrhea and predictors of its pain intensity among Palestinian female university students. *BMC Womens Health* 2018;18:18.
- 56 Hailemeskel S, Demissie A, Assefa N: Primary dysmenorrhea magnitude, associated risk factors, and its effect on academic performance: evidence from female university students in Ethiopia. *Int J Womens Health* 2016;8:489–496.
- 57 Pejčić A, Janković S: Risk factors for dysmenorrhea among young adult female university students. *Ann Ist Super Sanita* 2016;52:98–103.
- 58 Shinde GR, Laddad M: Overview of adolescent menstrual problems and its relation to BMI, eating habits and physical activity. *J Evolution Med Dental Sci* 2016;5:6757–6761.
- 59 Karacın O, Mutlu I, Kose M, Celik F, Kanat-Pektas M, Yilmazer M: Serum vitamin D concentrations in young Turkish women with primary dysmenorrhea: a randomized controlled study. *Taiwan J Obstet Gynecol* 2018; 57:58–63.
- 60 Muluneh AA, Nigussie TS, Gebreslasie KZ, Anteneh KT, Kassa ZY: Prevalence and associated factors of dysmenorrhea among secondary and preparatory school students in Debremarkos town, North-West Ethiopia. *BMC Womens Health* 2018;18:57.
- 61 Fujiwara T: Diet during adolescence is a trigger for subsequent development of dysmenorrhea in young women. *Int J Food Sci Nutr* 2007;58:437–444.
- 62 Mehrpooya M, Eshraghi A, Rabiee S, Larki-Harchegani A, Ataei S: Comparison the effect of fish-oil and calcium supplementation on treatment of primary dysmenorrhea. *Rev Recent Clin Trials* 2017;12:148–153.
- 63 Sadeghi N, Paknezhad F, Rashidi Nooshabadi M, Kavianpour M, Jafari Rad S, Khadem Haghighian H: Vitamin E and fish oil, separately or in combination, on treatment of primary dysmenorrhea: a double-blind, randomized clinical trial. *Gynecol Endocrinol* 2018; 34:804–808.
- 64 Zafari M, Tofighi M, Aghamohammady A, Behmanesh F, Rakhshaei Z: Comparison of the effect of acupuncture, fish oil capsules and ibuprofen on treatment of primary dysmenorrhea. *Afr J Pharm Pharmacol* 2011;5:1115–1119.
- 65 Saini RK, Nile SH, Park SW: Carotenoids from fruits and vegetables: chemistry, analysis, occurrence, bioavailability and biological activities. *Food Res Int.* 2015;76:735–750.
- 66 Moini A, Ebrahimi T, Shirzad N, Hosseini R, Radfar M, Bandarian F, et al: The effect of vitamin D on primary dysmenorrhea with vitamin D deficiency: a randomized double-blind controlled clinical trial. *Gynecol Endocrinol* 2016;32:502–505.
- 67 Zarei S, Mohammad-Alizadeh-Charandabi S, Mirghafourvand M, Javadzadeh Y, Effati-Daryani F: Effects of calcium-vitamin D and calcium-alone on pain intensity and menstrual blood loss in women with primary dysmenorrhea: a randomized controlled trial. *Pain Med* 2017;18:3–13.